

FLIPS AND VARIATION OF MODULI SCHEME OF SHEAVES ON A SURFACE

KIMIKO YAMADA

ABSTRACT. Let H be an ample line bundle on a non-singular projective surface X , and $M(H)$ the coarse moduli scheme of rank-two H -semistable sheaves with fixed Chern classes on X . We show that if H changes and passes through walls to get closer to K_X , then $M(H)$ undergoes natural flips with respect to canonical divisors. When X is minimal and $\kappa(X) \geq 1$, this sequence of flips terminates in $M(H_X)$; H_X is an ample line bundle lying so closely to K_X that the canonical divisor of $M(H_X)$ is nef. Remark that so-called Thaddeus-type flips somewhat differ from flips with respect to canonical divisors.

1. INTRODUCTION

Let X be a non-singular projective surface over $\mathbb{C}$, and H an ample line bundle on X . Denote by $M(H)$ (resp. $\bar{M}(H)$) the coarse moduli scheme of rank-two H -stable (resp. H -semistable) sheaves on X with Chern class $(c_1, c_2) \in \text{Pic}(X) \times \mathbb{Z}$. We shall consider birational aspects of the problem how $\bar{M}(H)$ changes as H varies.

There is a union of hyperplanes $W \subset \text{Amp}(X)$ called (c_1, c_2) -walls in the ample cone $\text{Amp}(X)$ such that $\bar{M}(H)$ changes only when H passes through walls. Let H_- and H_+ be ample line bundles separated by just one wall W . For $a \in (0, 1)$ one can define a -semistability of sheaves and the coarse moduli scheme $M(a)$ (resp. $\bar{M}(a)$) of rank-two a -stable (resp. a -semistable) sheaves with Chern classes (c_1, c_2) in such a way that $\bar{M}(\epsilon) = \bar{M}(H_+)$ and $\bar{M}(1 - \epsilon) = \bar{M}(H_-)$ if $\epsilon > 0$ is sufficiently small. Let $a_- < a_+$ be parameters separated by only one miniwall a_0 ; see Section 2 about miniwalls. Denote $\bar{M}_\pm = \bar{M}(a_\pm)$ and $\bar{M}_0 = \bar{M}(a_0)$. There are natural morphisms $f_- : \bar{M}_- \rightarrow \bar{M}_0$ and $f_+ : \bar{M}_+ \rightarrow \bar{M}_0$. After [4], let $f : X \rightarrow Y$ be a birational proper morphism such that K_X is $\mathbb{Q}$ -Cartier and $-K_X$ is f -ample, and that the codimension of the exceptional set $\text{Ex}(f)$ of f is more than 2. We say a birational proper morphism $f_+ : X_+ \rightarrow Y$ is a *flip* of f if (1) K_{X_+} is $\mathbb{Q}$ -Cartier, (2) K_{X_+} is f_+ -ample and (3) the codimension of the exceptional set $\text{Ex}(f_+)$ is more than 2. We do not require that each contraction reduces the Picard number by 1. The main result is the following.

Theorem 1.1. *Assume c_2 is so large that $\bar{M}_-$ and $\bar{M}_+$ are normal and that the codimensions of*

$$(1) \quad \bar{M}_\pm \supset P_\pm = \{[E] \mid E \text{ is not } a_\mp\text{-semistable}\}$$

and


$$\bar{M}_\pm \supset \text{Sing}(\bar{M}_\pm) = \{[E] \mid \dim \text{Ext}^2(E, E)^0 \neq 0\}$$

1991 *Mathematics Subject Classification.* Primary 14J60; Secondary 14E05, 14D20.
Revised on Dec. 12, 2008.

are more than 2. Suppose K_X does not lie in the wall W separating H_- and H_+ , and that K_X and H_- lie in the same connected components of $\text{NS}(X)_{\mathbb{R}} \setminus W$. (See the left figure below.) Then the birational map

$$(2) \quad \begin{array}{ccc} M_+ & \xrightarrow{\quad \quad \quad} & M_- \\ & \searrow f_- & \swarrow f_+ \\ & f_-(M_-) = f_+(M_+) \subset \bar{M}_0, & \end{array}$$

where $f_-(M_-)$ is open in $\bar{M}_0$, is a flip.



Let us observe this theorem in case where X is minimal and $\kappa(X) \geq 1$. When X is of general type, K_X is nef and big so there is an ample line bundle, say H_X , such that no wall of type (c_1, c_2) divides K_X and H_X from the proof of [6, Lem. 1.5]. Also when X is elliptic one can find such H_X from [2, p. 201]. Suppose $M(H) = \bar{M}(H)$ for any generic polarization H , that is valid if $c_1 = 0$ and c_2 is odd for example. Fix a compact polyhedral cone $\mathcal{S}$ in the closure of $\text{Amp}(X)$ with $\mathcal{S} \cap \partial \text{Amp}(X) \subset \mathbb{R}K_X$. If c_2 is sufficiently large, then $M(H)$ are isomorphic in codimension one for $H \in \mathcal{S}$. Hence when H starts from a polarization H_0 and gets closer to K_X , one gets flips

$$M(H) = M(a_0 = \epsilon) \cdots > M(a_1) \cdots > M(a_{N'} = 1 - \epsilon) = M(H')$$

when H passes through a wall to get closer to K_X by Theorem 1.1, where a_i are parameters lying in adjacent miniwalls. Hence the birational map $M(H) \cdots > M(H')$ is decomposed into finite sequence of natural flips. After we repeat it finitely many times, H reaches K_X and hence the sequence of birational morphisms

$$M(H = H_0) \cdots > M(H_1) \cdots > M(H_N = H_X)$$

terminates in $M(K_X)$. (See the right figure above.) It is known that the canonical divisor of $M(H_X)$ is nef. Thus one can regard this “natural” process described in a moduli-theoretic way as an analogy of minimal model program of $M(H)$, although it is unknown whether $M(K_X)$ admits only terminal singularities. Note that $M(H)$ is of general type if X is of general type, $H^0(K_X)$ contains a reduced curve, and $\chi(\mathcal{O}_X) + c_1^2(E) \equiv 0(2)$ by [5].

We mention some characteristics of this paper compared with Thaddeus’ work [8], which carefully and widely considered the variation of GIT quotients and linearizations. First of all, the argument here goes independently. As pointed out in [6], the rational map $M_- \rightarrow M_+$ is a Thaddeus-flip, that is, a rational map which is an isomorphism in codimension 1 and comes from the variation of GIT quotient and linearization. Thaddeus showed that the rational map $X//G(-) \cdots > X//G(+)$ is a flip with respect to $\mathcal{O}(1) \rightarrow X//G(+)$ at [8, Theorem 3.3]. Generally the relation about K -flip, that is, flip with respect to the canonical divisor is not mentioned there. So-called Thaddeus-flip is weaker than the flip defined above. Moreover the

birational map (2) is described concretely. Indeed, birational maps $M_{\pm} \rightarrow M_0$ associate a sheaf $E \in P_{\pm}$ with the graded sheaves (F, G) of the H_0 -Jordan-Hölder filtration

$$0 \longrightarrow F \longrightarrow E \longrightarrow G \longrightarrow 0;$$

see [1, Prop. 3.14]. Moduli schemes M_- and M_+ are connected by a natural blow-up and a blow-down described in moduli theory; see [9, Prop. 4.9].

2. PROOF OF THEOREM

There is a union of hyperplanes $W \subset \text{Amp}(X)$ called (c_1, c_2) -walls in the ample cone $\text{Amp}(X)$ such that $\bar{M}(H)$ changes only when H passes through walls ([7]). Let H_- and H_+ be ample line bundles separated by just one wall W , and $H_0 = \lambda H_- + (1 - \lambda)H_+$ an ample line bundle contained in W . Let H_- and H_+ be ample line bundles separated by just one wall W . If c_2 is sufficiently large with respect to a compact subset $S \subset \text{Amp}(X)$ containing $H_{\pm}$, then $M_{\pm}$ are normal and the codimension of $P_{\pm} \subset M_{\pm}$, which is defined at (1), are greater than 2 from [5] and [3, Thm. 4.C.7]. By [1, Sect. 3], for a number $a \in [0, 1]$ one can define the a -stability of a torsion-free sheaf E using

$$P_a(E(n)) = \{(1 - a)\chi(E(H_-)(nH_0)) + a\chi(E(H_+)(nH_0))\} / \text{rk}(E).$$

There is the coarse moduli scheme $\bar{M}(a)$ of rank-two a -semistable sheaves on X with Chern classes (c_1, c_2) . Denote by $M(a)$ its open subscheme of a -stable sheaves. There is a finite numbers a_i called miniwall such that, as a varies, $M(a)$ changes only when a passes through miniwalls. Let $a_- < a_+$ be parameters separated by only one miniwall a_0 , and denote $\bar{M}_{\pm} = \bar{M}(a_{\pm})$ and $\bar{M}_0 = \bar{M}(a_0)$. Since a rank-two a_- -semistable sheaf of type (c_1, c_2) is a_0 -semistable, there are natural morphisms $f_- : \bar{M}_- \rightarrow \bar{M}_0$ and $f_+ : \bar{M}_+ \rightarrow \bar{M}_0$ ([1, Prop. 3.14]).

Remark that the canonical divisors of $M_{\pm}$ are $\mathbb{Q}$ -Cartier. Indeed, M_- equal $R // \text{SL}(N)$, where R is a subscheme of the Quot-scheme parameterizing quotient sheaves $\mathcal{O}_X(-M) \rightarrow E^-$ on X . Let E_R^- be the universal quotient sheaf on X_R . From descent lemma [3, Theorem 4.2.15], $\det R\text{Hom}_{X_R/R}(E_R^-, E_R^-)$ descends to a line bundle on M_- , which we denote by $\det R\text{Hom}_{X_{M_-}/M_-}(E^-, E^-)$. It is known that $K_{M_-}|_{M_- \setminus \text{Sing}(M_-)}$ equals $\det R\text{Hom}_{X_{M_-}/M_-}(E^-, E^-)$ from deformation theory. Since M_- is normal, we have

$$(3) \quad K_{M_-} = \det R\text{Hom}_{X_{M_-}/M_-}(E^-, E^-),$$

so it is $\mathbb{Q}$ -Cartier.

Let η be an element of

$$A^+(W) = \{\eta \in \text{NS}(X) \mid \eta \text{ defines } W, 4c_2 - c_1^2 + \eta^2 \geq 0 \text{ and } \eta \cdot H_+ > 0\}.$$

After [1, Definition 4.2] we define

$$T_{\eta} = M(1, (c_1 + \eta)/2, n) \times M(1, (c_1 - \eta)/2, m),$$

where n and m are numbers defined by

$$n + m = c_2 - (c_1 - \eta^2)/4 \quad \text{and} \quad n - m = \eta \cdot (c_1 - K_X)/2 + (2a - 1)\eta \cdot M,$$

and $M(1, (c_1 + \eta)/2)$ is the moduli scheme of rank-one torsion-free sheaves on X with Chern classes $((c_1 + \eta)/2, n)$. If $F_{T_{\eta}}$ (resp. $G_{T_{\eta}}$) is the pull-back of a universal

sheaf of $M(1, (c_1 + \eta)/2, n)$ (resp. $M(1, (c_1 - \eta)/2, m)$) to X_{T_η} , then we have the following.

Proposition 2.1 ([9], Section 5). *We have isomorphisms*

$$(4) \quad P_- \simeq \coprod_{\eta \in A^+(W)} \mathbf{P}_{T_\eta} \left(\text{Ext}_{X_{T_\eta}/T_\eta}^1(F_{T_\eta}, G_{T_\eta}(K_X)) \right) \quad \text{and}$$

$$(5) \quad P_+ \simeq \coprod_{\eta \in A^+(W)} \mathbf{P}_{T_\eta} \left(\text{Ext}_{X_{T_\eta}/T_\eta}^1(G_{T_\eta}, F_{T_\eta}(K_X)) \right).$$

There are line bundles L_i (resp. L'_i) on P_- (resp. P_+) with exact sequences

$$\begin{aligned} 0 \longrightarrow F_T \otimes L_1 \longrightarrow E_{M_-}^-|_{P_-} \longrightarrow G_T \otimes L_2 \longrightarrow 0 \quad \text{and} \\ 0 \longrightarrow G_T \otimes L'_1 \longrightarrow E_{M_+}^+|_{P_+} \longrightarrow F_T \otimes L'_2 \longrightarrow 0 \end{aligned}$$

such that $L_1 \otimes L_2^{-1} = \mathcal{O}_{P_-}(1)$, which means the tautological line bundle of the right side of (4), and $L'_1 \otimes L'^{-1}_2 = \mathcal{O}_{P_+}(1)$. Here $E_{M_-}^-$ is a universal family of M_- , which exists etale-locally.

Claim 2.2. It holds that

$$\begin{aligned} K_{M_-}|_{P_- \times_T T_\eta} &= -(\eta \cdot K_X) \mathcal{O}_{P_-}(1) + (\text{some line bundle on } T), \quad \text{and} \\ K_{M_+}|_{P_+ \times_T T_\eta} &= (\eta \cdot K_X) \mathcal{O}_{P_+}(1) + (\text{some line bundle on } T). \end{aligned}$$

Proof. Suppose that $A^+(W) = \{\eta\}$ for simplicity. From the definition of walls, one can check $c_1(F_t) - c_1(G_t) = \eta$. By Proposition 2.1 and (3),

$$\begin{aligned} K_{M_-}|_{P_-} &= \det R\text{Hom}_{X_{P_-}/P_-}(E_{M_-}^-|_{P_-}, E_{M_-}^-|_{P_-}) = \\ &= \det R\text{Hom}_{X_T/T}(F_T, F_T) + \det(R\text{Hom}_{X_T/T}(F_T, G_T) \otimes L_2 \otimes L_1^{-1}) \\ &+ \det(R\text{Hom}_{X_T/T}(G_T, F_T) \otimes L_1 \otimes L_2^{-1}) + \det R\text{Hom}_{X_T/T}(G_T, G_T) \\ &= -\chi(F_t, G_t) \cdot \mathcal{O}_{P_-}(1) + \chi(G_t, F_t) \cdot \mathcal{O}_{P_-}(1) + (\text{line bundle on } T) \\ &= -(\eta \cdot K_X) \mathcal{O}_{P_-}(1) + (\text{line bundle on } T). \end{aligned}$$

One can calculate $K_{M_+}|_{P_+}$ similarly. $\square$

Remark that, since $\eta \cdot H_+ > 0$, one can verify that $\eta \cdot K_X < 0$ if and only if K_X does not lie in $W^\eta = W$, and H_- and K_X lie in the same connected components of $\text{NS}(X)_\mathbb{R} \setminus W$. Therefore we obtain Theorem 1.1.

We end with proving some facts in Introduction.

Lemma 2.3. *Fix a compact polyhedral cone $\mathcal{S}$ in the closure of $\text{Amp}(X)$ with $\mathcal{S} \cap \partial \text{Amp}(X) \subset \mathbb{R}K_X$. If c_2 is sufficiently large, then for $H \in \mathcal{S}$ $M(H)$ are isomorphic in codimension one.*

Proof. Remark that, for any $L \in \{tK_X + (1-t)H_0 \mid 0 \leq t < 1\}$,

$$|(K_X \cdot L)^2/L^2| \leq (K_X \cdot H_0)^2/H_0^2.$$

One can check this lemma in a similar way to [7, Thm. 2.3]. $\square$

Claim 2.4. The canonical bundle of $M(H_X)$ is nef.

Proof. From [3, Prop. 8.3.1] $2K_{M(H_X)} = p_*(\Delta(E_{M(H_X)}) \cdot K_X)$, and $p_*(\Delta(E_{M(H_f)}) \cdot H_X)$ is nef. $M(H) = M(H_X)$ if a polarization H is sufficiently close to K_X , so the claim follows. $\square$

REFERENCES

- [1] G. Ellingsrud and L. Göttsche, *Variation of moduli spaces and Donaldson invariants under change of polarization*, J. Reine Angew. Math. **467** (1995), 1–49.
- [2] R. Friedman, *Algebraic surfaces and holomorphic vector bundles*, Springer-Verlag, New York, 1998.
- [3] D. Huybrechts and M. Lehn, *The geometry of moduli spaces of sheaves*, Friedr. Vieweg & Sohn, 1997.
- [4] J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, vol. 134, Cambridge University Press, 1998.
- [5] J. Li, *Kodaira dimension of moduli space of vector bundles on surfaces*, Invent. Math. **115** (1994), no. 1, 1–40.
- [6] K. Matsuki and R. Wentworth, *Mumford-Thaddeus principle on the moduli space of vector bundles on an algebraic surface*, Internat. J. Math. **8** (1997), no. 1, 97–148.
- [7] Z. Qin, *Birational properties of moduli spaces of stable locally free rank-2 sheaves on algebraic surfaces*, Manuscripta Math. **72** (1991), no. 2, 163–180.
- [8] M. Thaddeus, *Geometric invariant theory and flips*, J. Amer. Math. Soc. **9** (1996), no. 3, 691–723.
- [9] K. Yamada, *A sequence of blowing-ups connecting moduli of sheaves and the Donaldson polynomial under change of polarization*, J. Math. Kyoto Univ. **43** (2003), no. 4, 829–878, math.AG/0704.2866.

E-mail address: kyamada@math.kyoto-u.ac.jp

DEPARTMENT OF MATHEMATICS, KYOTO UNIVERSITY, JAPAN